

REDUCING UNNECESSARY MEDICINES

Taking fewer medicines can have major benefits for both health and the environment. It reduces the risk of drug interactions and side effects while also limiting pharmaceutical residues that contaminate ecosystems, particularly rivers and lakes.¹ By questioning each prescription, we can improve quality of life and care while also helping to protect the environment.

34%

of Swiss insured persons over 65 years received a potentially inappropriate prescription in 2019.²

32 000

hospitalizations per year in Switzerland are due to adverse drug reactions.³

60 tons

of medications have been found in Lake Geneva, Switzerland.



The **environmental impact of medication production and consumption** is equivalent to that of land transport, based on evaluations in some European countries (France, UK).

How to reduce medication use ?

The example of non-steroidal anti-inflammatory drugs (NSAIDs) commonly used to reduce fever, inflammation, and pain.

GOOD TO KNOW

To relieve pain, **non-drug alternatives** exist: physiotherapy, hot/cold applications, osteopathy, acupuncture, and (self-)hypnosis.⁴



BENEFITS FOR HEALTH

Reducing NSAID use helps prevent their side effects: **stomach pain, cardiovascular issues, and kidney problems.**⁵

NSAIDs interact with many medications (e.g., **anticoagulants, antidepressants**), either enhancing or reducing their effects.⁵ Reducing their use helps **avoid these undesirable interactions.**



BENEFITS FOR THE ENVIRONMENT

Reducing NSAID consumption **helps protect wildlife**, since these medicines can harm the livers of fish and contribute to the decline of birds of prey by affecting their kidneys.¹

Lower medication consumption reduces **water pollution and prevents waste.**¹

THE SUSTAINABLE PRESCRIPTION

- ◆ Avoid unnecessary prescriptions.
- ◆ Reevaluate each prescription.
- ◆ Consider non-drug alternatives.
- ◆ For physicians: Use Smarter Medicine recommendations (www.smartermedicine.ch) and START/STOP criteria.⁶

When to discuss de-prescription or NSAID reduction ?

During **each prescription renewal or when introducing a new medication.**
For NSAIDs: particularly in cases of **chronic and/or acute pain**, during **consultations for flu-like symptoms or viral infections.**

REFERENCES

1. Senn N, Gaillie M, del Río Carral M, Gonzalez Holguera J (dirs). Health and Environment: Toward a New Global Approach. RMS Edition, 2022. Free PDF download.

2. Vallon N, et al. Use of Potentially Inappropriate Medications in the Canton of Valais and Inter-canton Comparison. Unisanté, Lausanne. Nov. 2023.

3. Beeler PE, et al. Hospitalizations Related to Adverse Drug Reactions in Switzerland (2012-2019): Characteristics, In-Hospital Mortality, and Spontaneous Reporting Rate. Drug Saf. 2023; 46(8): 753-63.

4. Alternative Pain Management Methods [CHUM [Internet]. [cited Oct 15, 2024]. Available at: <https://www.chumontreal.qc.ca/fiches-sante/autres-moyens-pour-gerer-douleur>

5. Bourdon F, et al. Non-Steroidal Anti-Inflammatory Drugs: A Practical Update. Rev Med Suisse. 2022; 18(768): 235-40.

6. Lang PO, et al. Inappropriate Medication Prescribing: The New STOPP/START Criteria. Rev Med Suisse. Nov 11, 2015; 494: 2115-23.



UNIVERSITÉ
DE GENÈVE
FACULTÉ DE MÉDECINE



Société
Genevoise
de Pédiatrie

REVUE
MÉDICALE
SUISSE

